

Abstracts

1. **Molnár I**, Kaczur V., Kovács I., Balázs C.: IgA autoantibodies against human eye muscle tissue in Graves' disease. (Abstract) *Period Biol* 1994; 96: 269.
2. **Molnár I**, Balázs C: Elevated IL-6 levels in Graves' ophthalmopathy. (Abstract) *J Endocrinol Invest* 1996; 19 (Suppl.No.6):83.
3. **Molnár I**, Szabó E, Balázs C: The presence of anti-TSH antibodies in the euthyroid state of Graves' disease. (Abstract) *J Endocrinol Invest* 1997; 20 (Suppl 5):89.
4. **Molnár I**, Stief L: Iodine deficiency in patients with cardiovascular disease. (Abstract) *Eur J Internal Med* 1997; 8 (Suppl 1):61.
5. **Molnár I**, Czirják L.: Changes of thyroid hormone levels in systemic sclerosis. (Abstract) *J Endocrinol Invest* 1998; 21 (Suppl. No.4): 60.
6. **Molnár I**, Sipka S, Szegedi Gy, Balázs Cs: Direct action of IL-6, TNF α and IFN γ on the activity of type 2 5'-deiodinase in human thyroid tissue. (Abstract) *J Endocrinol Invest* 1999; 22 (Suppl. No 6):36.
7. Csathy L, **Molnár I**: Iodine determinations in newborn infants. (Abstract) *Arch Hung Med Assoc Am* 1999; 7:2.
8. **Molnár I**, Balázs C: Study of serum IL-5 and IL-12 levels in Graves' disease. (Abstract) *ACI International* 2000; Suppl. 2:220.
9. **Molnár I**, Csathy L: Autoantibodies against the type 2 5'-deiodinase are involved in the development of Graves' ophthalmopathy. (Abstract) *Endocrinol Japonica* 2000; 47 (Suppl.): 176.
10. Csathy L, **Molnár I**, Szasz G, Math J: Determination of thyroid gland volume by ultrasound and iodine excretion in newborns in East Hungary. (Abstract) *Endocrinol Japonica* 2000; 47 (Suppl.):236.
11. **Molnár I**, Balázs Cs.: Elevated serum levels of IL-12 in Graves' ophthalmopathy. (Abstract) *J Endocrinol Invest* 2001; 24 (Suppl.6):86.
12. **Molnár I**, Szombathy Z, Kovacs I, Szentmiklosi J: Detection of deiodinase in human thyroid and skeletal muscle tissues using sera of guinea pigs immunized against corresponding to human type 2 5'-deiodinase. Abstract Book, 73rd Annual Meeting of the American Thyroid Association, (Abstract) Washington, USA, 2001, p 123.
13. **Molnár I**, Csathy L, Szasz G: Relationship of gestational age with the neonatal urinary iodine excretion and the thyroid volume. Abstract Book, 73rd Annual Meeting of the American Thyroid Association, (Abstract) Washington, USA, 2001, p 176.
14. **Molnár I**, Csathy L, Szasz G, Math J: Thyroid gland volume, iodine excretion and type of feeding in newborns. Abstract Book, 73rd Annual Meeting of the American Thyroid Association, (Abstract) Washington, USA, 2001, p 238.

15. **Molnár I**, Bokk Á: Serum levels of nerve growth factor in autoimmune and nonimmune thyroid diseases. (Abstract) *J Endocrinol Invest* 2002; 25 (Suppl. 7):90.
16. **Molnár I**, Szentmiklósi J : The activation of mitogen-activated protein kinase (MAPK) in lymphocytes of patients with autoimmune thyroid diseases. (Abstract). *Autoimmunity Reviews* 2004; 3 (Suppl 2):103.
17. **Molnár I**, Bokk Á : Graves'hyperthyroidism associated with elevated levels of nerve growth factor. (Abstract) *Exp Clin Endocrinol Diab* 2005; 113 (8):501.
18. **Molnár I**, Kelemen E : Characteristics of allergy in autoimmune thyroid diseases. (Abstract) *Thyroid* 2005; 15 (Suppl. 1):S-52.
19. Csáthy L, **Molnár I**, Kukla E, Orvos H, Vida G, Alexy M : Urinary iodine determination in newborns in Hungary. (Abstract) *Thyroid* 2005; 15 (Suppl. 1): S-182.
20. **Molnár I**, Bokk Á: Graves' hyperthyroidism associated with elevated levels of nerve growth factor. *Exp Clin Endocrinol Diabetes* 2005; 113:501.
21. **Molnar I**, Kelemen E: Relationship between age and allergens in patients with thyroid diseases. Abstract Book, Annual Meeting of the Austrian Society for Allergology and Immunology, (Abstract), Graz, Austria, 2005, p 92.
22. **Molnár I**, Kelemen E: Relationship between allergen sensitization and immuno-endocrine events in Graves' disease. Abstract Book, XXV Congress of the European Academy of Allergology and Clinical Immunology (Abstract) Vienna, Austria 2006, p 431.
23. **Molnár I**, Láng A: The role of IL-1 receptor antagonists in the eye symptoms of Graves' ophthalmopathy. Book of Abstracts, 16th European Congress of Immunology, (1st Join Meeting of European National Societies of Immunology) (Abstract), Paris, France, 2006, p 572.
24. **Molnár I**: Sensitization against Soybean may induce an increase in the levels of anti-thyroid peroxidase antibodies in thyroid autoimmunity. *Endocrine Abstracts*, 9th European Congress of Endocrinology (Abstract), Budapest, Hungary 2007; Vol 14, OC1.6.
25. **Molnar I**, Kelemen E: Allergen-specific IgE may contribute to the change of thyroid hormones in thyroid autoimmunity. Abstract Book, Annual Meeting of the Austrian Society for Allergology and Immunology, (Abstract), Vienna, Austria 2010, A-191-OEGAI.
26. **Molnar I**, Bokk Á: The regulatory role of neuroprotective nerve growth factor in Graves' ophthalmopathy. *Endocrine Abstracts*, 13th European Congress of Endocrinology, (Abstract), Rotterdam, The Netherlands, 2011; Vol 26, P423.
27. **Molnar I**, Kelemen E, Somogyine-Vari E: Antigen mimicry between aeroallergens and thyroid antigens can modify the levels of thyroid hormones and antibodies in thyroid autoimmunity. *Endocrine Abstracts*, 15th International & 14th European Congress of Endocrinology (ICE/ECE 2012) (Abstract), Florence, Italy 2012; Vol 29, P1606.

28. **Molnár I**, Bohaty I, Somogyiné-Vári É: Increased IL-17A serum levels associated with low estrogen levels can play a role in postmenopausal osteoporosis. Endocrine abstracts & 15th European Congress of Endocrinology (Abstract), Copenhagen, Denmark, 2013; Vol 32: P80.
29. Bohaty I, Jurásné-Lukovics M, Somogyiné-Vári É, Kozma L, Dankó K, **Molnár I**: HLA class I, Cw*01 and Cw*15 alleles can play a preventing role in serum IL-17 elevation associated with postmenopausal osteoporosis in Hungary. Endocrine abstracts & 15th European Congress of Endocrinology (Abstract), Copenhagen, Denmark, 2013; Vol 32: P85.
30. **Molnár I**, Bohaty I, Somogyiné-Vári É: 1,25-dihydroxyvitamin D₃ decreases the pro-inflammatory interleukin 17 serum levels. Endocrine abstracts & 16th European Congress of Endocrinology (Abstract), Wrocław, Poland, 2014; Vol 35: P80.
31. **Molnár I**, Timkó T, Somogyiné-Vári É: Increased occurrence of proinflammatory IL-17, IL-6 and MCP-1 cytokines and resistin with the overweight-associated polycystic ovarian syndrome, pituitary adenome and obesity. Endocrine abstracts & 16th European Congress of Endocrinology (Abstract), Wrocław, Poland, 2014; Vol 35: P633.
32. **Molnár I**, Bohaty I, Somogyiné-Vári É: Smoking associated with TSH receptor antibody and serum IL-17A elevation in hyperthyroid Graves' disease. Endocrine abstracts & 17th European Congress of Endocrinology (Abstract), Dublin, Ireland 2015; Vol 37: EP934.
33. **Molnár I**, Szentmiklósi AJ, Somogyiné-Vári É: Effect of patient sera on type 2 5'-deiodinase enzyme activity using human thyroid, skeletal and eye muscle tissue fractions in hyperthyroid Graves' ophthalmopathy. Abstract Book, 15th International Thyroid Congress & 85th Annual Meeting of the American Thyroid Association (Abstract), Orlando, USA, 2015, P515.
34. **Molnár I**, Szentmiklósi AJ, Somogyiné-Vári É: Antibodies against human eye muscle tissue fractions bind to type 2 5'-deiodinase and inhibit the lymphocyte MAPK activation derived from hyperthyroid patients with Graves' disease. Endocrine abstracts & 18th European Congress of Endocrinology (Abstract), Munich 2016; Vol 41: OC4.3.
35. **Molnár I**, Somogyiné-Vári É: Distinct tissue-specific effects of methylprednisolone and pentoxifylline on type 2 5'-deiodinase enzyme activities using thyroid and eye muscle tissue fractions. Abstract Book, 86th Annual Meeting of the American Thyroid Association, (Abstract), Denver, USA, 2016, P122.
36. MIT 46th Annual Meeting of the Hungarian Society for Immunology, 18-20th October 2017, Velence
Molnár I, Szentmiklósi AJ, Gesztelyi R, Somogyiné-Vári É: Influencing effect of medications on the frequency of antibodies against protein sequences of type 2 deiodinase and their relationships with the thyroid hormone and antibody levels in hyperthyroid Graves' ophthalmopathy